

Genetic and Genomics aspects of susceptibility and resistance to infections

Cesar Paz-y-Miño¹✉, Javier Gualdron², Ariane Paz-y-Miño³

¹ Equinoccial Technological University - Faculty of Health Sciences «Eugenio Espejo», Ecuador

² National blood center of the Ecuadorian Red Cross-Laboratory of Molecular Biology, Ecuador

³ Mass General Brigham Salem Hospital, United States

Competing interests: CPM none; JG none; APM none

Abstract

This article addresses the genetic and genomic determinants that influence susceptibility or resistance to human infections, whether caused by bacteria, viruses, fungi, parasites or even prions. A large number of genes are involved in susceptibility or resistance, and this is because *Homo sapiens sapiens* evolutionarily acquired key DNA modifications adaptable to defense as a result of the interaction of infectious agents with their genetic properties, natural selection and mutations. This article provides a description of the methods used in the investigation of this evolution and mutual host-infectious agent adaptation. The genes involved in susceptibility and resistance are presented, as well as the polymorphic variants that provide greater or lesser reaction capacity. The high number of genes, which produce specific defense proteins, have intricate relationships among them. Relationships among these genes have been analysed. A list of variants, and even mutations of the 70 to 153 proteins/genes that are involved in the immune response to infection diseases has been compiled. Additionally, we designed a network of physical interactions as predictable, with BioGrid data and enrichment data obtained by the g:Profiler platform, finding 57 of the 70 genes with protein-to-protein interactions. The network detected is composed of 1,049 interacting genes (in total 1,106 genes and 1,910 interactions). This analysis shows the importance of immunity genes in the defense against infectious agents, as well as the effects of the genes involved such as *HLA*, immunoglobulins, interleukins, immune cells, among others. Based on the functional analysis of genes involved in susceptibility and resistance to infections, we compiled a list of genetic disorders that increase an individual's risk of developing infectious diseases. It is known that infectious diseases manifest general clinical characteristics, which are a good guide to suspect a disease. Furthermore, we describe general clinical signs that may suggest underlying genetic immunodeficiency, and highlight specific infectious diseases—such as HIV/AIDS, amebiasis, malaria, Chagas disease, tuberculosis, and COVID-19—where genetic susceptibility factors have been well characterized.

Introduction

Infectious diseases continue to be one of the leading causes of morbidity and mortality worldwide and the variability in susceptibility or resistance of individuals to these infections highlights the role of genetic factors. The ability of the immune system to respond to infectious agents such as viruses, bacteria, fungi, parasites and even prions is influenced by genetic variants, which may predispose to develop infections or, conversely, confer resistance to these (Bos *et al.*, 2019; Quintana-Murci, 2019).

The genetics of human immunity has been shaped throughout evolution, in a process of natural selection and adaptation to diverse environments. This process

has resulted in key modifications to the DNA of *H. sapiens*, allowing specific genetic variants to optimise the immune defense against infectious agents (Quintana-Murci, 2019). In addition, the environment in which individuals live, such as tropical or temperate zones and interactions with other species have influenced the genetic diversity of immunity. Thus, coevolution with pathogens has imposed significant selective pressure on certain genes, favoring the permanence of variants that increase survival in contexts of high infectious exposure (Vasseur and Quintana-Murci, 2013; Quintana-Murci, 2019).

Several genes influence susceptibility and resistance to AIDS-causing HIV. The absence of TNPO3 confers

Article history

Received: 24 April 2025

Accepted: 06 July 2025

Published: September 2025

© 2025 Paz-y-Miño *et al.*; the authors have retained copyright and granted the Journal right of first publication; the work has been simultaneously released under a Creative Commons Attribution Licence, which allows others to share the work, while acknowledging the original authorship and initial publication in this Journal. The full licence notice is available at <http://journal.embnet.org>.

absolute resistance, while other genes such as CPSF6 and SC35 are also involved. Mutations in CCR5, CXCR4 and SDF1 receptors increase susceptibility in homozygosis. In Ecuador, resistance mutations are present in 0.5% (CCR5), 16% (CCR2) and 48% (SDF1), but only in heterozygosis, which does not guarantee total resistance (Paz-y-Miño Cepeda *et al.*, 2005). In Europe, genetic resistance is 26%, while in Latin America it is only 4% (Chapman and Hill, 2012; McLaren *et al.*, 2015).

Amoeba infections are a public health problem, influenced by innate and adaptive immunity genes. Mutations in IL8 and TNF- α increase the risk of *Entamoeba histolytica* infection; variations in MHC affect susceptibility, since this gene regulates the immune response. On the other hand, inflammatory genes such as IL1 β and TLR2 are also associated with increased risk and mutations in epithelial barrier genes, such as filaggrin and TLR4, favor infection (Haque *et al.*, 2002; Guo *et al.*, 2011; Robledo *et al.*, 2018). Whether the host genome influences the localization of infection, along with parasite factors, is debated.

Malaria, caused by *Plasmodium* and transmitted by *Anopheles*, is influenced by genetic and environmental factors. In Ecuador, 1,946 cases were reported in 2020, mainly due to *P. vivax* (Muñoz Cabas *et al.*, 2023). The mutation in the HBB gene (HbS) confers resistance to *P. falciparum* in heterozygotes, but increases the risk in homozygotes (Hill *et al.*, 1991; Malaria Genomic Epidemiology Network *et al.*, 2019; Muñoz Cabas *et al.*, 2023). In Esmeraldas, 24.3% of the population has hemoglobin variants, with Hb.AS as the most common, other mutations in HBC, HBE and G6PD also influence resistance, with a G6PD prevalence of 10% in endemic areas. *Plasmodium vivax*, is found in 11% of the Afro-Ecuadorian population and genes such as TLR1, IL10 and PfEMP1, together with MHC and TNE, also affect susceptibility and severity of malaria (Auburn and Barry, 2017; Muñoz Cabas *et al.*, 2023).

Chagas disease, caused by *Trypanosoma cruzi*, affects about 8 million people worldwide. In Ecuador, 113 cases were reported in 2020, mostly chronic (Dumonteil *et al.*, 2016). Genetic factors influence susceptibility and severity, including immune system genes, MHC, IL-10, IL-6 and SOD, which affect the inflammatory response and oxidative stress (Dumonteil *et al.*, 2016). Variants in POLD1 and repair genes may contribute to cell damage and a study of SNPs found no association with cardiomyopathy, but identified 44 SNPs associated with various disease traits (Vasconcelos *et al.*, 2012; Deng *et al.*, 2013; Frade *et al.*, 2013).

Tuberculosis is a multifactorial disease influenced by genetic and environmental factors. In Ecuador, in 2020, the incidence was 24 cases per 100,000 inhabitants, with an increase of 34% in deaths and 9.4% in cases (Anon, 2018). Genes such as NRAMP1 and IFNG are associated with increased susceptibility, affecting the immune response (McNicholl *et al.*, 2000; Davila *et al.*, 2008). TNF- α , TLRs, CISH and TYK2 also play a role, the latter with variants that have reduced incidence in

the last 4,000 years, and VDR, IL10, IL12A, IL12B, IL6, IL17A and IL17F impact immunity and risk of pulmonary tuberculosis (Hawn *et al.*, 2007; African TB Genetics Consortium *et al.*, 2010; Cholo *et al.*, 2015; Curtis *et al.*, 2015).

Susceptibility to COVID-19 is influenced by genetic factors such as blood type, ethnicity, autoimmunity genes, HLA, and heart or kidney failure genes. The ACE2 receptor, located on chromosome Xp22.2, is key in the entry of SARS-CoV-2 into lung cells, likewise, a study identified 45 proteins associated with susceptibility, highlighting 11 with strong interactions, such as ACE2, AGT, AGTR1, REN and DPP4, related to blood pressure regulation and hormone metabolism (Debnath *et al.*, 2020; Paz-y-Miño *et al.*, 2021; Gupta *et al.*, 2022; Saengsiwaritt *et al.*, 2022; Pecoraro *et al.*, 2023). These interactions explain the clinical heterogeneity of the disease and its impact on different body systems.

The study of the genetic determinants associated with the immune response to infections not only provides a basis for a better understanding of the pathogenesis of these diseases, but also opens the possibility of developing personalised therapies. Several genes and their variants are associated with susceptibility or resistance to specific infections. For example, major histocompatibility complex (HLA) genes, Toll-like receptors (TLRs) and cytokines play an essential role in the innate and adaptive response, affecting the immune system's ability to recognize and destroy pathogens (Merker *et al.*, 2020; Martins *et al.*, 2023; Paz y Miño Cepeda, 2024).

This study analyzes the key genes involved in the immune response to infections and describes how the construction of protein-protein interaction networks, or interactomes, helps to understand the functional relationships between them. By analyzing these networks, we identify genes that act as "hubs" or central nodes in the immune response, revealing their role in resistance or susceptibility to infections such as HIV, tuberculosis, malaria, and COVID-19 (Casadevall and Pirofski, 2000; Rast and Litman, 2010).

Materials and methods

Gene Selection

To identify genes related to resistance and susceptibility to infections we considered studies describing genes and genetic variants associated with changes in gene expression, alterations in RNA structure, modifications in immunoglobulin conformation, modulation of the immune response, abnormal interactions with other genes and the influence of environmental factors (Janeway and Medzhitov, 2002; Khor and Hibberd, 2012; Rolland *et al.*, 2014; Mozzi *et al.*, 2018; Luck *et al.*, 2020; Nahon and Cobat, 2020).

A systematic review of scientific literature was carried out in databases such as PubMed and Scopus, using key terms such as immunity genes, genetic susceptibility and infections. The selection criteria

included: (i) functional association with infections, (ii) evidence of relevant genetic polymorphisms, and (iii) involvement in immune processes. The 70 selected genes were categorised into innate, adaptive, inflammatory and antiviral immune responses. Table S1a¹ presents some groups of genes relevant in this context, giving an overview of their contribution to resistance and susceptibility to infections, while Table S2a² presents the genes selected for this study.

To solve the problem related to genes codifying for different proteins but having the same names, we have consulted GeneCards ([genecard.com](http://www.genecard.com)) (Stelzer *et al.*, 2016). Here it is possible to locate and designate each protein with its proper gene using its accession number in the UniProt³ database (The UniProt Consortium *et al.*, 2023; in Supplementary Statistical Material, Table S1⁴).

Construction of Protein-Protein Interaction (PPI) Networks

Experimental and predicted data from BioGrid (60) and STRING (Oughtred *et al.*, 2021; Szklarczyk *et al.*, 2023) were used to map protein-protein interactions. PPI networks were generated using the Cytoscape software (Doncheva *et al.*, 2019), following the steps:

- Nodes represented genes/proteins and edges indicated physical or functional interactions.
- The MCODE algorithm was used to identify functional clusters in the network.
- Interactions were filtered for reliability according to the scores provided by STRING.

Functional Enrichment

Functional analysis was performed with Gprofiler (Kolberg *et al.*, 2023), considering statistically significant terms ($p < 0.05$) from Gene Ontology (GO) (in Supplementary Statistical Material, Table S3⁵) (64), KEGG (in Supplementary Statistical Material, Table S4⁶) (65) and Reactome (in Supplementary Statistical Material, Table S5⁷) (Ashburner *et al.*, 2000; Kanehisa, 2000; Rothfels *et al.*, 2023). Pathways related to biological

¹http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_S1a

²http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_S2a

³www.uniprot.org

⁴http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_1

⁵http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_1

⁶http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_1

⁷http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_1

processes, such as cytokine activation, immune signaling and inflammatory regulation were evaluated. Enriched terms were visualised using Manhattan diagrams generated in Cytoscape (Doncheva *et al.*, 2019).

Topological Analysis

Key topological metrics of PPI networks were analysed to identify critical nodes (“hubs”) using Cytoscape (in Supplementary Statistical Material, Table S2⁸) (Doncheva *et al.*, 2019).

- Degree of connection: nodes with the highest number of interactions.
- Intermediation centrality: nodes that acted as control points between interactions.
- Identification of key genes such as TNF, HBB and CD79B, prioritised for their impact on immunity.

Results and discussions

Susceptibility- and resistance-related proteins listed in Table S2a⁹ were subjected to a bioinformatics analysis of protein-protein interactions (PPIs), defined as highly specific physical contacts between two or more proteins, driven by electrostatic forces, hydrogen bonds, or hydrophobic effects. These interactions were used to construct a protein interaction network, or interactome (Figure 1) (Rolland *et al.*, 2014; Luck *et al.*, 2020). The interactome was generated using the STRING database, and may serve as a reference model for understanding complex protein interactions involved in the immune response to infections (Szklarczyk *et al.*, 2023).

Related pathologies

The PPI network in Figure 1 reveals that among all the proteins analysed ($n=70$) there are 624 interactions, with a total correlation score of 0.669. Such correlation indicating that the proteins are at least partially biologically connected, as a group. Each protein involved in the interactome has a known function and these functions are observed in different pathological manifestations (Stelzer *et al.*, 2016; Szklarczyk *et al.*, 2023; The UniProt Consortium *et al.*, 2023). Table 1 shows the diseases related to their anomalous interactions, therefore, product of mutations or genetic variants, resulting from the analysis of the proteins/genes interactome produced.

Additionally, the ontological analysis carried out by the String Software’s of the 70 genes analysed, yields at least 17 functional nodes congruent with different functional pathways such as: Immune receptor activation, transmembrane signaling receptor activations, virus receptor activation, 2-5-oligoadenylate synthetase activation, activation of regulators of molecular

⁸http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_1

⁹http://journal.embnet.org/index.php/embnetjournal/article/downloadSuppFile/1073/1073_supp_S2a

Table 1. Diseases related to abnormal protein-gene interactions and infection predisposition genes

Disease	Monogenic (M) and/or Polygenic (P) Component*	Disease	Monogenic (M) and/or Polygenic (P) Component*
X-linked Hyper IgM Syndrome	M	Guillain-Barré Syndrome	M, P
Immunodeficiency with Hyper IgM Type 3	M	Genital Herpes	P
Common Cold	P	Cutaneous Leishmaniasis	P
Pulmonary Eosinophilia	P	Visceral Leishmaniasis	P
Adult Respiratory Distress Syndrome	P	Hyperimmunoglobulin Syndrome	M
Stevens-Johnson Syndrome	M	Mantle Cell Lymphoma	M, P
Rosacea	P	Acquired Immune Deficiency Syndrome	P
Influenza	P	Conjunctivitis	P
Membranous Glomerulonephritis	P	Lymphocytopenia	P
Intrinsic Asthma	P	Leprosy	P
Behçet's Disease	M	COVID-19	P
Allergic Asthma	P	Allergic Rhinitis	M, P
Myocarditis	P	Encephalomyelitis	P
Crohn's Disease	P	Vasculitis	P
Infectious Disease Due to Parasites and Protozoa	P	B-cell Deficiency	M
Glomerulonephritis	P	Leukopenia	P
Hypersensitivity Disease and Type IV Reaction	P	Viral Infections	P
Pneumonia	P	Parasitic Infections	P
Helminth Disease	P	Asthma	P
Bronchial Disease	P	Leukocyte Disease	M, P
Primary Bacterial Infection	P	Liver Disease	M, P
Infectious Agent Disease	P	Primary Immunodeficiency	M
Upper Respiratory Tract Disease	P	Pulmonary Disease	P
Bacterial Infections	P	Dermatitis	P
Allergic Diseases	M, P	Renal Disease	M, P
Lower Respiratory Tract Diseases	P	Skin Diseases	M, P
Immune System Diseases	M, P	Vascular Diseases	M, P
Urinary System Diseases	M, P	Carcinoma	M, P
Intestinal Diseases	M, P	Various Genetic Diseases	M
Connective System Diseases	M	Human Immunodeficiency Virus	M, P
Diseases of the Gastrointestinal System	M, P		

*Each disease listed is associated with either monogenic (M) or polygenic (P) mutations that influence susceptibility to infections. These genetic variations can impact disease severity and symptomatology. For example, the CCR5 gene provides resistance to HIV infection in homozygous individuals but not in heterozygous ones, illustrating a monogenic effect. However, other genes also contribute to susceptibility or resistance, demonstrating the polygenic nature of many conditions.

all the metrics yielded by the analysis, we used the degree of interaction of the nodes to define the “centers” or Hubs of connection in the network and the “Intermediation Centrality” as a measure of the frequency with which a node appears in the shortest paths between other nodes (Figure 2 B). High values in this metric define critical control points or bottlenecks in the network (Wu *et al.*, 2008; Koh *et al.*, 2012). We found that CD79B, TNF, TYK2, POLD1, BCR and HBB genes are the “hubs” with the highest degree of connection in the analysed network. The identification of these “hub nodes” underlines their possible roles in immune response and pathogen defense mechanisms. CD79B and BCR are integral parts of the B-cell receptor complex and influence adaptive immune responses (Pleiman *et al.*, 1994; Tanaka and Baba, 2020), while TNF is a fundamental cytokine involved

in systemic inflammation (Bulló *et al.*, 2003), crucial for infection control, but also implicated in inflammatory diseases (Rickert, 2013). TYK2, part of the JAK-STAT signaling pathway, is key to the signaling of several type I interferons and cytokines, influencing both innate and adaptive immunity (Villarino *et al.*, 2017). POLD1, although primarily associated with DNA replication and repair (Nicolas *et al.*, 2016), may have functions in the immune system that are less studied, but potentially significant, particularly in terms of genomic stability in rapidly proliferating immune cells (Nichols-Vinueza *et al.*, 2021). Finally, HBB, part of the hemoglobin complex, is critical in oxygen transport, but also plays a role in modulating the response to oxidative stress during infections (Ma, 2013; Ramezani *et al.*, 2018). These “hubs”, being highly interconnected within the




Figure 2. The topological analysis of the protein interaction network.

Note* Enrichment analysis and protein-protein interaction network associated with susceptibility to infection. A) Manhattan-type graphical representation showing the most significant ontology terms derived from the enrichment analysis. Terms are color-coded according to their origin: molecular function terms from the Gene Ontology (GO) in red, signaling pathways from the KEGG database in green and pathways from the REACTOME database in purple. B) Protein-protein interaction network constructed from BioGrid database interaction data. The topology of the network highlights the central nodes (red boxes) according to their degree of interaction with the proteins studied and their association with different cytokines (purple), clusters of differentiation (CDs) (red), major histocompatibility complex (HLA) molecules (green), and Toll-like receptors (TLRs) (yellow). The size of the nodes reflects the "Centrality of Intermediation" index within the network.

network, suggest that they not only play individual roles in the response to infection, but may also interact synergistically, offering potential strategic targets for improving resistance to infection and understanding mechanisms of susceptibility.

Immune cells, such as lymphocytes, macrophages and dendritic cells, detect and destroy invading pathogens. Immune system proteins, such as immunoglobulins, interferons and cytokines, help regulate and coordinate the immune response. Failures

in the genetic construction of these elements are associated with the development of disease (Janeway and Medzhitov, 2002; Khor and Hibberd, 2012; Mozzi *et al.*, 2018; Rotival, 2019; Nahon and Cobat, 2020). Interestingly, protein interaction network analysis significantly weights cytokines, human leukocyte antigen (HLA) genes and Toll-like receptors (TLRs) as genes associated with fundamental mechanisms of resistance or susceptibility to infection (Szklarczyk *et al.*, 2023). Each group of proteins contributes uniquely to the

immune system's ability to defend against pathogens are presented in Table S3a¹¹.

Cytokine are signaling molecules that mediate and regulate immunity, inflammation and hematopoiesis (Kany *et al.*, 2019). Genetic variants in this group of proteins, such as in the gene encoding the interleukin-23 (IL-23) receptor, have been linked to susceptibility to inflammatory bowel disease and fungal infections. Likewise, genetic variants in interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF- α) are associated with increased disease severity in patients with COVID-19 (Khor and Hibberd, 2012; Mozzi *et al.*, 2018; Horwood *et al.*, 2019; Masin *et al.*, 2022). The presence of IFNAR1, IFNAR2, IL10, IL4, IL4R, IL6ST and TNF in our interaction network indicates their role in orchestrating immune responses. Cytokines such as TNF and IL6 participate in the acute inflammatory response, which is crucial for controlling the spread of infection (Rankin, 2004). In contrast, regulatory cytokines such as IL10 and IL4 can modulate the immune response to prevent over activation, which can lead to tissue damage or autoimmune disorders (Couper *et al.*, 2008; Junttila, 2018). The balance in cytokine signaling may dictate the outcome of an infection, so the study of the aforementioned proteins may improve the description of effective pathogen clearance processes or susceptibility due to an uncontrolled inflammatory response.

Major histocompatibility complex (MHC) proteins, also known as the human leukocyte antigen (HLA) group, play a crucial role in the immune response by facilitating the recognition of foreign molecules by the immune system. These proteins are essential for the presentation of antigens to T cells, enabling the identification and elimination of pathogens (Chaplin, 2010). In the context of Lyme disease for example, a tick-bite-borne infection caused by the bacterium *Borrelia burgdorferi*, certain genetic variants in the HLA genes, specifically HLA-DRB1, HLA-DQA1 and HLA-DQB1, have been found to be linked to an increased susceptibility to this disease (Mozzi *et al.*, 2018; Rotival, 2019). Molecular variants in these genes significantly influence the host's ability to respond to infection, highlighting the importance of genetic polymorphisms in the pathogenesis of infectious diseases.

On the other hand, Toll-like receptors (TLRs) are fundamental components of the innate immune system, acting as essential molecular sensors in the detection of pathogens. These receptors identify highly conserved structures called pathogen-associated molecular patterns (PAMPs), characteristics of invading microorganisms. When activated, TLRs initiate signaling cascades that elicit inflammatory responses and trigger crucial defense mechanisms, including cytokines production and effector cells activation (Kumar *et al.*, 2009; Kawai and Akira, 2010, 2011). This rapid and widespread responsiveness is vital for immediate

defense against infection and facilitates the transition between innate and adaptive immunity by influencing the activation and maturation of cells of the adaptive immune system (O'Neill *et al.*, 2013). Genetic variations in the genes encoding for TLRs can significantly modify the efficacy of the innate immune system in recognising and responding appropriately to PAMPs (Vijay, 2018). Such variations can result in attenuated or over-activated immune responses, increasing susceptibility to infections, as in the case of tuberculosis or predisposing to chronic inflammatory responses (Khor and Hibberd, 2012; Walker *et al.*, 2015; Mozzi *et al.*, 2018; Paz-y-Miño *et al.*, 2021). Our network analysis identifies TLR1, TLR2, TLR3, TLR4 and TLR7 as central nodes in pathogen susceptibility. These TLRs are crucial not only for the recognition of pathogen-associated molecular patterns, but also for triggering innate immune responses and subsequent modular adaptive responses. Their central position in the network underscores their pivotal role in the initial detection of and response to infectious agents (Janeway and Medzhitov, 2002; Khor and Hibberd, 2012; Masin *et al.*, 2022).

Weighing these groups in network analysis provides deep insight into the collective and individual contributions to immune system functionality, highlighting how imbalances or deficiencies in any of these components could lead to increased susceptibility to infections, while optimal functioning and interaction confer resistance. Understanding these dynamics is essential for developing interventions that aim to stimulate immune responses and effectively manage infection-related diseases.

Identification of disease clusters

Central clusters within the network were identified using Mcode and possible disease associations with the analysed network were explored. The analysis of clusters and diseases with high significance in the analysed network is presented in Table 2.

Many of the diseases presented in Table 2 are caused by microorganisms, and genetic variability among them can also influence their aggressiveness or resistance to antimicrobials. In the adaptive game of evolution, infectious agents do their best to escape the immune system (selective pressure), and to do so they frequently change their genomes (Khor and Hibberd, 2012; Mozzi *et al.*, 2018; Masin *et al.*, 2022). This is precisely the reason for the difficulty of effective treatments, new resistances and the impossibility of designing vaccines.

Immune cells, such as lymphocytes, macrophages and dendritic cells, detect and destroy invading pathogens. Immune system proteins, such as immunoglobulins, interferons and cytokines, help regulate and coordinate the immune response. Failures in the genetic construction of these elements are associated with the development of disease (Janeway and Medzhitov, 2002; Khor and Hibberd, 2012; Mozzi *et al.*, 2018). Genetic and functional determinants of defence

¹¹http://journal.embnet.org/index.php/embnetjournal/article/download/SuppFile/1073/1073_supp_S3a

Table 2. Diseases associated with gene network clusters.

GO/KEGG Term	Adjusted p-value
Hepatitis B	1.14E-30
Epstein-Barr virus infection	2.35E-29
Lipids and atherosclerosis	5.92E-26
Measles	2.94E-24
Kaposi's sarcoma-associated herpesvirus infection	9.60E-22
Influenza A	5.77E-21
Hepatitis C	9.77E-21
Shigellosis	1.48E-19
Toxoplasmosis	4.14E-19
Human cytomegalovirus infection	9.70E-19
Yersinia infection	1.20E-17
Viral carcinogenesis	1.96E-16
Tuberculosis	5.02E-16
Human T-cell leukemia virus infection	6.80E-16
Chronic myeloid leukemia	2.21E-15
Human immunodeficiency virus infection	9.08E-15
Coronavirus disease - COVID-19	1.47E-14
Salmonella infection	1.73E-13
Human papillomavirus infection	4.66E-12
Necroptosis	9.04E-12
Chagas disease	9.31E-12
Alcoholic liver disease	3.38E-10
Pancreatic cancer	5.44E-10
Leishmaniasis	8.10E-10
Shear stress and atherosclerosis	2.57E-09
Prostate cancer	2.85E-09
Small cell lung cancer	3.40E-09
Renal cell carcinoma	4.82E-05
Malaria	2,56E-01
Acute myeloid leukemia	3,69E-01
Glioma	6,18E-01
Non-small cell lung cancer	0.00013277

mechanisms allow most people to recover fully from infections, although there are cases in which an infection can have serious or even fatal consequences. The groups of genes with a role in the resistance or susceptibility to infections are presented in Table S1a¹².

The genetic variants involved in the impairment of the immune response to infections include changes in the immunoglobulin chain configuration, B cell behavior and elevation, HLA system organisation, participation in the inflammatory response, complement system activities, cytokine, receptor and enzymes involved in immunoglobulin maturation molecular conformation (Janeway and Medzhitov, 2002; Khor and Hibberd, 2012; Rotival, 2019).

Susceptibility to infectious diseases may be due to mutations in genes encoding defense proteins. Some clear examples such as mutations in the gene encoding the interleukin-12 receptor, interleukin-1 and variants in

¹²http://journal.embnet.org/index.php/embnetjournal/article/download/SuppFile/1073/1073_supp_S1a

the TLR2 gene, which encodes the Toll-like receptor 2, have been associated with susceptibility to tuberculosis infection in at-risk populations (McNicholl *et al.*, 2000; Hawn *et al.*, 2007; Davila *et al.*, 2008; Curtis *et al.*, 2015). The gene encoding the interleukin-23 receptor (IL23R) is implicated in susceptibility to inflammatory bowel disease and fungal infections. Additionally, IL23R is a cytokine that stimulates the production of inflammatory cytokines and activates immune cells, which help fight fungal infections (Junttila, 2018; Kany *et al.*, 2019).

The FUT2 gene, which encodes an enzyme involved in the synthesis of cell surface antigens, is associated with increased susceptibility to norovirus and rotavirus infections. Individuals with mutations in the FUT2 gene cannot produce certain antigens that fight these viral infections (Mozzi *et al.*, 2018; Masin *et al.*, 2022).

The IFITM3 gene encodes a protein that helps prevent entry of the influenza virus into cells, individuals who carry one copy of a gene variant have an increased risk of infection and hospitalisation for influenza (Mozzi *et al.*, 2018; Horwood *et al.*, 2019; Rasch, 2019).

In Lyme disease caused by the tick-borne bacterium *Borrelia burgdorferi*, genetic variants in the HLA-DRB1, HLA-DQA1 and HLA-DQB1 genes, which code for major histocompatibility complex (MHC) molecules, are known to be associated with increased susceptibility to this disease (De Los Rios *et al.*, 2015; Mozzi *et al.*, 2018).

Genetic variants in the genes encoding for the cytokines IL6 and TNF- α are associated with increased disease severity in patients with COVID-19, so identification of these genetic variants may help to better understand the pathogenesis of the disease and develop new therapies (Khor and Hibberd, 2012; Walker *et al.*, 2015; Mozzi *et al.*, 2018; Debnath *et al.*, 2020; Masin *et al.*, 2022).

Evolutionarily, genetic variation involves the suitable or unsuitable production of immunoglobulins, resulting in greater antibody diversity, greater or lesser affinity and specificity of these antibodies and individual immune response (Bachman and LeBar, 2018; Nielsen and Boyd, 2018). Table S4a¹³ shows some diseases of genetic origin and their international classification (OMIM) (Amberger and Hamosh, 2017), which confer greater predisposition to infections.

The common symptoms and signs that raise suspicion of a genetic deficiency or variant predisposing to infections, or a genetic disease with a risk of infections, are shown in Table S5a¹⁴ (Nielsen and Boyd, 2018; Carroll and Pfaller, 2023).

Conclusions

This study analysed the genetic basis of human susceptibility and resistance to infections by identifying

¹³http://journal.embnet.org/index.php/embnetjournal/article/download/SuppFile/1073/1073_supp_S4a

¹⁴http://journal.embnet.org/index.php/embnetjournal/article/download/SuppFile/1073/1073_supp_S5a

key genes, their polymorphic variants, and the protein–protein interaction networks shaping the immune interactome. The aim was to understand how genetic architecture influences immune defense mechanisms and to provide novel interpretative approaches through bioinformatic and topological analyses.

Notably, we constructed an extensive interaction network comprising 1,106 genes and 1,910 interactions, identifying central hub genes such as CD79B, TNE, TYK2, POLD1, and HBB. These genes, although known individually, had not previously been jointly prioritised as immune control nodes in the literature. Moreover, the integration of databases such as BioGrid and g:Profiler enabled functional enrichment analysis, highlighting immune-related pathways including cytokine signaling, Toll-like receptor activation, and the MHC complex.

These findings provide a comprehensive framework for future investigations into the molecular mechanisms of immune response to infection. They may contribute to the development of personalised therapies, identification of genetic susceptibility biomarkers, and a better understanding of infectious diseases through the lens of evolutionary biology and precision medicine.

The dataset and prioritised genes proposed here may serve as a valuable resource for translational and clinical research in immunogenetics.

Key Points

- Human susceptibility or resistance to infections is strongly influenced by genetic variants that modulate immune response.
- An immune interactome was constructed with 1,106 genes and 1,910 interactions, identifying key hub genes such as TNE, CD79B, and TYK2.
- Functional pathway analysis revealed enrichment in immune-related processes including cytokine activation, TLR signaling, and the MHC complex.
- Specific genetic variants are associated with diseases such as HIV/AIDS, tuberculosis, malaria, and COVID-19.
- Understanding these genetic interactions provides valuable insights into personalised medicine in the context of infectious diseases.

References

- African TB Genetics Consortium, The Wellcome Trust Case Control Consortium, Thyne T, Vannberg FO, Wong SH, Owusu-Dabo E, et al. Genome-wide association analyses identifies a susceptibility locus for tuberculosis on chromosome 18q11.2. *Nat Genet.* 2010;42:739–741. <http://dx.doi.org/10.1038/ng.639>.
- Akbay B, Germini D, Bissenbaev AK, Musinova YR, Sheval EV, Vassetzky Y, et al. HIV-1 Tat Activates Akt/mTORC1 Pathway and AICDA Expression by Downregulating Its Transcriptional Inhibitors in B Cells. *Int J Mol Sci.* 2021;22:1588. <http://dx.doi.org/10.3390/ijms22041588>.
- Amato KR, Jeyakumar T, Poinar H, Gros P. Shifting Climates, Foods, and Diseases: The Human Microbiome through Evolution. *BioEssays.* 2019;41:1900034. <http://dx.doi.org/10.1002/bies.201900034>.
- Amberger JS, Hamosh A. Searching Online Mendelian Inheritance in Man (OMIM): A Knowledgebase of Human Genes and Genetic Phenotypes. *Curr Protoc Bioinforma [Internet].* 2017 [cited 2024 Nov 5];58. <http://dx.doi.org/10.1002/cpbi.27>.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology: tool for the unification of biology. *Nat Genet.* 2000;25:25–29. <http://dx.doi.org/10.1038/75556>.
- Auburn S, Barry AE. Dissecting malaria biology and epidemiology using population genetics and genomics. *Int J Parasitol.* 2017;47:77–85. <http://dx.doi.org/10.1016/j.ijpara.2016.08.006>.
- Bachman MA, LeBar WD. First Principles of Clinical Microbiology: Collection, Handling, and Diagnostics. Ref Module Biomed Sci [Internet]. Elsevier; 2018 [cited 2024 Nov 5]. p. B9780128012383661160. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780128012383661160>.
- Baranova A, Cao H, Chen J, Zhang F. Causal Association and Shared Genetics Between Asthma and COVID-19. *Front Immunol.* 2022;13:705379. <http://dx.doi.org/10.3389/fimmu.2022.705379>.
- Baranova A, Cao H, Zhang F. Unraveling Risk Genes of COVID-19 by Multi-Omics Integrative Analyses. *Front Med.* 2021;8:738687. <http://dx.doi.org/10.3389/fmed.2021.738687>.
- Baxter VK, Heise MT. Immunopathogenesis of alphaviruses. *Adv Virus Res [Internet].* Elsevier; 2020 [cited 2024 Dec 16]. p. 315–382. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0065352720300208>.
- Beigier-Bompadre M, Montagna GN, Kühl AA, Lozza L, Weiner J, Kupz A, et al. Mycobacterium tuberculosis infection modulates adipose tissue biology. Behr MA, editor. *PLOS Pathog.* 2017;13:e1006676. <http://dx.doi.org/10.1371/journal.ppat.1006676>.
- Boletín anual de Tuberculosis [Internet]. Ministerio de Salud Pública del Ecuador; 2018. Available from: [www: https://www.salud.gov.ec/gacetitas-tuberculosis/](http://www.salud.gov.ec/gacetitas-tuberculosis/).
- Bos KI, Kühnert D, Herbig A, Esquivel-Gomez LR, Andrades Valtueña A, Barquera R, et al. Paleomicrobiology: Diagnosis and Evolution of Ancient Pathogens. *Annu Rev Microbiol.* 2019;73:639–666. <http://dx.doi.org/10.1146/annurev-micro-090817-062436>.
- Bulló M, García-Lorda P, Megias I, Salas-Salvadó J. Systemic Inflammation, Adipose Tissue Tumor Necrosis Factor, and Leptin Expression. *Obes Res.* 2003;11:525–531. <http://dx.doi.org/10.1038/oby.2003.74>.
- Butler-Laporte G, Kreuzer D, Nakanishi T, Harroud A, Forgetta V, Richards JB. Genetic Determinants of Antibody-Mediated Immune Responses to Infectious Diseases Agents: A Genome-Wide and HLA Association Study. *Open Forum Infect Dis.* 2020;7:ofaa450. <http://dx.doi.org/10.1093/ofid/ofaa450>.
- Carroll KC, Pfaller MA, editors. Manual of clinical microbiology. Washington, DC: ASM Press; 2023.
- Casadevall A, Pirofski L. Host-Pathogen Interactions: Basic Concepts of Microbial Commensalism, Colonization, Infection, and Disease. Portnoy DA, editor. *Infect Immun.* 2000;68:6511–6518. <http://dx.doi.org/10.1128/IAI.68.12.6511-6518.2000>.
- Chaplin DD. Overview of the immune response. *J Allergy Clin Immunol.* 2010;125:S3–S23. <http://dx.doi.org/10.1016/j.jaci.2009.12.980>.
- Chapman SJ, Hill AVS. Human genetic susceptibility to infectious disease. *Nat Rev Genet.* 2012;13:175–188. <http://dx.doi.org/10.1038/nrg3114>.
- Chen L, Hua J, He X. Bioinformatics analysis identifies a key gene HLA_DPA1 in severe influenza-associated immune infiltration. *BMC Genomics.* 2024;25:257. <http://dx.doi.org/10.1186/s12864-024-10184-7>.
- Chen L-Y, Hwang A-C, Huang C-Y, Chen L-K, Wang F-D, Chan Y-J. CMV infection, CD19+ B cell depletion, and Lymphopenia as predictors for unexpected admission in the institutionalized elderly. *Immun Ageing.* 2021;18:21. <http://dx.doi.org/10.1186/s12979-021-00233-0>.
- Chen Z, Chen R, Ou Y, Lu J, Jiang Q, Liu G, et al. Construction of an HLA Classifier for Early Diagnosis, Prognosis, and Recognition of Immunosuppression in Sepsis by Multiple Transcriptome Datasets. *Front Physiol.* 2022;13:870657. <http://dx.doi.org/10.3389/fphys.2022.870657>.

- Choi H, Lee JY, Yoo H, Jeon K. Bioinformatics Analysis of Gene Expression Profiles for Diagnosing Sepsis and Risk Prediction in Patients with Sepsis. *Int J Mol Sci.* 2023;24:9362. <http://dx.doi.org/10.3390/ijms24119362>.
- Cholo MC, Van Rensburg EJ, Osman AG, Anderson R. Expression of the Genes Encoding the Trk and Kdp Potassium Transport Systems of *Mycobacterium tuberculosis* during Growth In Vitro. *BioMed Res Int.* 2015;2015:1–11. <http://dx.doi.org/10.1155/2015/608682>.
- Costafreda MI, Kaplan G. HAVCR1 (CD365) and Its Mouse Ortholog Are Functional Hepatitis A Virus (HAV) Cellular Receptors That Mediate HAV Infection. Pfeiffer JK, editor. *J Virol.* 2018;92:e02065-17. <http://dx.doi.org/10.1128/JVI.02065-17>.
- Couper KN, Blount DG, Riley EM. IL-10: The Master Regulator of Immunity to Infection. *J Immunol.* 2008;180:5771–5777. <http://dx.doi.org/10.4049/jimmunol.180.9.5771>.
- Cui Y, Keles S, Charbonnier L-M, Julé AM, Henderson L, Celik SC, et al. Combined immunodeficiency caused by a loss-of-function mutation in DNA polymerase delta 1. *J Allergy Clin Immunol.* 2020;145:391–401.e8. <http://dx.doi.org/10.1016/j.jaci.2019.10.004>.
- Curtis J, Luo Y, Zenner HL, Cuchet-Lourenço D, Wu C, Lo K, et al. Susceptibility to tuberculosis is associated with variants in the ASAP1 gene encoding a regulator of dendritic cell migration. *Nat Genet.* 2015;47:523–527. <http://dx.doi.org/10.1038/ng.3248>.
- Da Silva FAR, Pascoal LB, Dotti I, Setsuko Ayriszono MDL, Aguilar D, Rodrigues BL, et al. Whole transcriptional analysis identifies markers of B, T and plasma cell signaling pathways in the mesenteric adipose tissue associated with Crohn's disease. *J Transl Med.* 2020;18:44. <http://dx.doi.org/10.1186/s12967-020-02220-3>.
- Davila S, Hibberd ML, Hari Dass R, Wong HEE, Sahiratmadja E, Bonnard C, et al. Genetic Association and Expression Studies Indicate a Role of Toll-Like Receptor 8 in Pulmonary Tuberculosis. Gojobori T, editor. *PLoS Genet.* 2008;4:e1000218. <http://dx.doi.org/10.1371/journal.pgen.1000218>.
- De Los Rios M, Criscitiello ME, Smider VV. Structural and genetic diversity in antibody repertoires from diverse species. *Curr Opin Struct Biol.* 2015;33:27–41. <http://dx.doi.org/10.1016/j.sbi.2015.06.002>.
- Debnath M, Banerjee M, Berk M. Genetic gateways to COVID-19 infection: Implications for risk, severity, and outcomes. *FASEB J.* 2020;34:8787–8795. <http://dx.doi.org/10.1096/fj.202001115R>.
- Deng X, Sabino EC, Cunha-Neto E, Ribeiro AL, Ianni B, Mady C, et al. Genome Wide Association Study (GWAS) of Chagas Cardiomyopathy in *Trypanosoma cruzi* Seropositive Subjects. Tanowitz HB, editor. *PLoS ONE.* 2013;8:e79629. <http://dx.doi.org/10.1371/journal.pone.0079629>.
- Desai JV, Kumar D, Freiwald T, Chauss D, Johnson MD, Abers MS, et al. C5a-licensed phagocytes drive sterilizing immunity during systemic fungal infection. *Cell.* 2023;186:2802–2822.e22. <http://dx.doi.org/10.1016/j.cell.2023.04.031>.
- Doncheva NT, Morris JH, Gorodkin J, Jensen LJ. Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data. *J Proteome Res.* 2019;18:623–632. <http://dx.doi.org/10.1021/acs.jproteome.8b00702>.
- Druszczynska M, Seweryn M, Wawrocki S, Kowalewska-Pietrzak M, Pankowska A, Rudnicka W. Cytokine Biosignature of Active and Latent *Mycobacterium tuberculosis* Infection in Children. *Pathogens.* 2021;10:517. <http://dx.doi.org/10.3390/pathogens10050517>.
- Duan T, Du Y, Xing C, Wang HY, Wang R-F. Toll-Like Receptor Signaling and Its Role in Cell-Mediated Immunity. *Front Immunol.* 2022;13:812774. <http://dx.doi.org/10.3389/fimmu.2022.812774>.
- Dumonteil E, Herrera C, Martini L, Grijalva MJ, Guevara AG, Costales JA, et al. Chagas Disease Has Not Been Controlled in Ecuador. Tanowitz HB, editor. *PLOS ONE.* 2016;11:e0158145. <http://dx.doi.org/10.1371/journal.pone.0158145>.
- Fan H, Lu B, Cao C, Li H, Yang D, Huang L, et al. Plasma TNFSF13B and TNFSF14 Function as Inflammatory Indicators of Severe Adenovirus Pneumonia in Pediatric Patients. *Front Immunol.* 2021;11:614781. <http://dx.doi.org/10.3389/fimmu.2020.614781>.
- Frade AF, Pissetti CW, Ianni BM, Saba B, Lin-Wang HT, Nogueira LG, et al. Genetic susceptibility to Chagas disease cardiomyopathy: involvement of several genes of the innate immunity and chemokine-dependent migration pathways. *BMC Infect Dis.* 2013;13:587. <http://dx.doi.org/10.1186/1471-2334-13-587>.
- Ghiglione Y, Falivene J, Ruiz MJ, Laufer N, Socías ME, Cahn P, et al. Early Skewed Distribution of Total and HIV-Specific CD8+ T-Cell Memory Phenotypes during Primary HIV Infection Is Related to Reduced Antiviral Activity and Faster Disease Progression. Boasso A, editor. *PLoS ONE.* 2014;9:e104235. <http://dx.doi.org/10.1371/journal.pone.0104235>.
- Giraldo MI, Xia H, Aguilera-Aguirre L, Hage A, Van Tol S, Shan C, et al. Envelope protein ubiquitination drives entry and pathogenesis of Zika virus. *Nature.* 2020;585:414–419. <http://dx.doi.org/10.1038/s41586-020-2457-8>.
- Gómez I, Thomas MC, Palacios G, Egui A, Carrilero B, Simón M, et al. Differential Expression of Immune Response Genes in Asymptomatic Chronic Chagas Disease Patients Versus Healthy Subjects. *Front Cell Infect Microbiol.* 2021;11:722984. <http://dx.doi.org/10.3389/fcimb.2021.722984>.
- Goretzki A, Zimmermann J, Lin Y-J, Schülke S. Immune Metabolism—An Opportunity to Better Understand Allergic Pathology and Improve Treatment of Allergic Diseases? *Front Allergy.* 2022;3:825931. <http://dx.doi.org/10.3389/falgy.2022.825931>.
- Grant M, Larijani M. Evasion of adaptive immunity by HIV through the action of host APOBEC3G/F enzymes. *AIDS Res Ther.* 2017;14:44. <http://dx.doi.org/10.1186/s12981-017-0173-8>.
- Guo X, Roberts MR, Becker SM, Podd B, Zhang Y, Chua SC, et al. Leptin signaling in intestinal epithelium mediates resistance to enteric infection by *Entamoeba histolytica*. *Mucosal Immunol.* 2011;4:294–303. <http://dx.doi.org/10.1038/mi.2010.76>.
- Gupta K, Kaur G, Pathak T, Banerjee I. Systematic review and meta-analysis of human genetic variants contributing to COVID-19 susceptibility and severity. *Gene.* 2022;844:146790. <http://dx.doi.org/10.1016/j.gene.2022.146790>.
- Haque R, Duggal P, Ali IM, Hossain MB, Mondal D, Sack RB, et al. Innate and Acquired Resistance to Amebiasis in Bangladeshi Children. *J Infect Dis.* 2002;186:547–552. <http://dx.doi.org/10.1086/341566>.
- Hartman ZC, Kiang A, Everett RS, Serra D, Yang XY, Clay TM, Amalfitano A. Adenovirus Infection Triggers a Rapid, MyD88-Regulated Transcriptome Response Critical to Acute-Phase and Adaptive Immune Responses In Vivo. *J Virol.* 2007;81:1796–1812. <http://dx.doi.org/10.1128/JVI.01936-06>.
- Hawn TR, Misch EA, Dunstan SJ, Thwaites GE, Lan NTN, Quy HT, et al. A common human TLR1 polymorphism regulates the innate immune response to lipopeptides. *Eur J Immunol.* 2007;37:2280–2289. <http://dx.doi.org/10.1002/eji.200737034>.
- Heuberger J, Trimpert J, Vladimirova D, Goosmann C, Lin M, Schmuck R, et al. Epithelial response to IFN- γ promotes SARS-CoV-2 infection. *EMBO Mol Med.* 2021;13:e13191. <http://dx.doi.org/10.15252/emmm.202013191>.
- Hill AVS, Allsopp CEM, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA, et al. Common West African HLA antigens are associated with protection from severe malaria. *Nature.* 1991;352:595–600. <http://dx.doi.org/10.1038/352595a0>.
- Horwood PF, Tarantola A, Goarant C, Matsui M, Klement E, Umezaki M, et al. Health Challenges of the Pacific Region: Insights From History, Geography, Social Determinants, Genetics, and the Microbiome. *Front Immunol.* 2019;10:2184. <http://dx.doi.org/10.3389/fimmu.2019.02184>.
- Ishibashi M, Morita N, Nomura-Kawaguchi C, Shimizu Y, Wakita T, Esumi M. CLEC4M-positive and CD81-negative Huh7 cells are not susceptible to JFH-1 HCVcc infection but mediate transinfection.

- Arch Virol. 2014;159:2949–2955. <http://dx.doi.org/10.1007/s00705-014-2150-z>.
- Ismailova A, White JH. Vitamin D, infections and immunity. *Rev Endocr Metab Disord*. 2022;23:265–277. <http://dx.doi.org/10.1007/s11154-021-09679-5>.
- Janeway CA, Medzhitov R. Innate Immune Recognition. *Annu Rev Immunol*. 2002;20:197–216. <http://dx.doi.org/10.1146/annurev.immunol.20.083001.084359>.
- Josyula JVN, Talari P, Pillai AKB, Mutheneni SR. Analysis of gene expression profile for identification of novel gene signatures during dengue infection. *Infect Med*. 2023;2:19–30. <http://dx.doi.org/10.1016/j.imj.2023.02.002>.
- Junttila IS. Tuning the Cytokine Responses: An Update on Interleukin (IL)-4 and IL-13 Receptor Complexes. *Front Immunol*. 2018;9:888. <http://dx.doi.org/10.3389/fimmu.2018.00888>.
- Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res*. 2000;28:27–30. <http://dx.doi.org/10.1093/nar/28.1.27>.
- Kany S, Vollrath JT, Relja B. Cytokines in Inflammatory Disease. *Int J Mol Sci*. 2019;20:6008. <http://dx.doi.org/10.3390/ijms20236008>.
- Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol*. 2010;11:373–384. <http://dx.doi.org/10.1038/ni.1863>.
- Kawai T, Akira S. Toll-like Receptors and Their Crosstalk with Other Innate Receptors in Infection and Immunity. *Immunity*. 2011;34:637–650. <http://dx.doi.org/10.1016/j.immuni.2011.05.006>.
- Khasnis S, Veenstra H, McClellan MJ, Ojeniyi O, Wood CD, West MJ. Regulation of B cell receptor signalling by Epstein–Barr virus nuclear antigens. *Biochem J*. 2022;479:2395–2417. <http://dx.doi.org/10.1042/BCJ20220417>.
- Khor CC, Hibberd ML. Host–pathogen interactions revealed by human genome-wide surveys. *Trends Genet*. 2012;28:233–243. <http://dx.doi.org/10.1016/j.tig.2012.02.001>.
- Kitada S, Kayama H, Okuzaki D, Koga R, Kobayashi M, Arima Y, et al. BATF2 inhibits immunopathological Th17 responses by suppressing IL23a expression during *Trypanosoma cruzi* infection. *J Exp Med*. 2017;214:1313–1331. <http://dx.doi.org/10.1084/jem.20161076>.
- Koh GCKW, Porras P, Aranda B, Hermjakob H, Orchard SE. Analyzing Protein–Protein Interaction Networks. *J Proteome Res*. 2012;11:2014–2031. <http://dx.doi.org/10.1021/pr201211w>.
- Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler—interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). *Nucleic Acids Res*. 2023;51:W207–W212. <http://dx.doi.org/10.1093/nar/gkad347>.
- Kulkarni DH, Starick M, Aponte Albuquerque R, Kulkarni HS. Local complement activation and modulation in mucosal immunity. *Mucosal Immunol*. 2024;17:739–751. <http://dx.doi.org/10.1016/j.mucimm.2024.05.006>.
- Kumar H, Kawai T, Akira S. Toll-like receptors and innate immunity. *Biochem Biophys Res Commun*. 2009;388:621–625. <http://dx.doi.org/10.1016/j.bbrc.2009.08.062>.
- Kwissa M, Nakaya HI, Onlamoon N, Wrammert J, Villinger F, Perng GC, et al. Dengue Virus Infection Induces Expansion of a CD14+CD16+ Monocyte Population that Stimulates Plasmablast Differentiation. *Cell Host Microbe*. 2014;16:115–127. <http://dx.doi.org/10.1016/j.chom.2014.06.001>.
- Lang B, Cao C, Zhao X, Wang Y, Cao Y, Zhou X, et al. Correction: Genomic alterations related to HPV infection status in a cohort of Chinese prostate cancer patients. *Eur J Med Res*. 2024;29:348. <http://dx.doi.org/10.1186/s40001-024-01935-z>.
- Li X, Yang Y, Zhou F, Zhang Y, Lu H, Jin Q, et al. SLC11A1 (NRAMP1) Polymorphisms and Tuberculosis Susceptibility: Updated Systematic Review and Meta-Analysis. Herrmann JL, editor. *PLoS ONE*. 2011;6:e15831. <http://dx.doi.org/10.1371/journal.pone.0015831>.
- Lio CT, Dhanda SK, Bose T. Cluster Analysis of Dry Eye Disease Models Based on Immune Cell Parameters – New Insight Into Therapeutic Perspective. *Front Immunol*. 2020;11:1930. <http://dx.doi.org/10.3389/fimmu.2020.01930>.
- Loterio RK, Zamboni DS, Newton HJ. Keeping the host alive – lessons from obligate intracellular bacterial pathogens. *Pathog Dis*. 2021;79:ftab052. <http://dx.doi.org/10.1093/femspd/ftab052>.
- Luck K, Kim D-K, Lambourne L, Spirohn K, Begg BE, Bian W, et al. A reference map of the human binary protein interactome. *Nature*. 2020;580:402–408. <http://dx.doi.org/10.1038/s41586-020-2188-x>.
- Ma H, Liu J, Li Z, Xiong H, Zhang Y, Song Y, et al. Expression profile analysis reveals hub genes that are associated with immune system dysregulation in primary myelofibrosis. *Hematology*. 2021;26:478–490. <http://dx.doi.org/10.1080/16078454.2021.1945237>.
- Ma Q. Role of Nrf2 in Oxidative Stress and Toxicity. *Annu Rev Pharmacol Toxicol*. 2013;53:401–426. <http://dx.doi.org/10.1146/annurev-pharmtox-011112-140320>.
- Malaria Genomic Epidemiology Network, Band G, Le QS, Clarke GM, Kivinen K, Hubbard C, et al. Insights into malaria susceptibility using genome-wide data on 17,000 individuals from Africa, Asia and Oceania. *Nat Commun*. 2019;10:5732. <http://dx.doi.org/10.1038/s41467-019-13480-z>.
- Malaria Genomic Epidemiology Network, Rockett KA, Clarke GM, Fitzpatrick K, Hubbard C, Jeffreys AE, et al. Reappraisal of known malaria resistance loci in a large multicenter study. *Nat Genet*. 2014;46:1197–1204. <http://dx.doi.org/10.1038/ng.3107>.
- Malhotra S, Yen JY, Honko AN, Garamszegi S, Caballero IS, Johnson JC, et al. Transcriptional Profiling of the Circulating Immune Response to Lassa Virus in an Aerosol Model of Exposure. Geisbert T, editor. *PLoS Negl Trop Dis*. 2013;7:e2171. <http://dx.doi.org/10.1371/journal.pntd.0002171>.
- Martins YC, Ribeiro-Gomes FL, Daniel-Ribeiro CT. A short history of innate immunity. *Mem Inst Oswaldo Cruz*. 2023;118:e230023. <http://dx.doi.org/10.1590/0074-02760230023>.
- Masin PS, Visentin HA, Elpidio LNS, Sell AM, Visentainer L, Lima Neto QAD, et al. Genetic polymorphisms of toll-like receptors in leprosy patients from southern Brazil. *Front Genet*. 2022;13:952219. <http://dx.doi.org/10.3389/fgene.2022.952219>.
- Mathew NR, Jayanthan JK, Smirnov IV, Robinson JL, Axelsson H, Nakka SS, et al. Single-cell BCR and transcriptome analysis after influenza infection reveals spatiotemporal dynamics of antigen-specific B cells. *Cell Rep*. 2021;35:109286. <http://dx.doi.org/10.1016/j.celrep.2021.109286>.
- McLaren PJ, Coulonges C, Bartha I, Lenz TL, Deutsch AJ, Bashirova A, et al. Polymorphisms of large effect explain the majority of the host genetic contribution to variation of HIV-1 virus load. *Proc Natl Acad Sci*. 2015;112:14658–14663. <http://dx.doi.org/10.1073/pnas.1514867112>.
- McNicholl JM, Downer MV, Udhayakumar V, Alper CA, Swerdlow DL. Host-Pathogen Interactions in Emerging and Re-emerging Infectious Diseases: A Genomic Perspective of Tuberculosis, Malaria, Human Immunodeficiency Virus Infection, Hepatitis B, and Cholera. *Annu Rev Public Health*. 2000;21:15–46. <http://dx.doi.org/10.1146/annurev.publhealth.21.1.15>.
- Merker M, Tueffers L, Vallier M, Groth EE, Sonnenkalb L, Unterwiesing D, et al. Evolutionary Approaches to Combat Antibiotic Resistance: Opportunities and Challenges for Precision Medicine. *Front Immunol*. 2020;11:1938. <http://dx.doi.org/10.3389/fimmu.2020.01938>.
- Moroney JB, Chupp DP, Xu Z, Zan H, Casali P. Epigenetics of the antibody and autoantibody response. *Curr Opin Immunol*. 2020;67:75–86. <http://dx.doi.org/10.1016/j.coi.2020.09.004>.
- Mozzi A, Pontremoli C, Sironi M. Genetic susceptibility to infectious diseases: Current status and future perspectives from genome-wide approaches. *Infect Genet Evol*. 2018;66:286–307. <http://dx.doi.org/10.1016/j.meegid.2017.09.028>.

- Muñoz Cabas D, Cando Caluña W, Cando Herrera F, Cando Herrera J. Epidemiología de la Malaria en Ecuador de 2018–2022. RECIAMUC. 2023;7:151–162. [http://dx.doi.org/https://doi.org/10.26820/reciamuc/7.\(1\).enero.2023.151-162](http://dx.doi.org/https://doi.org/10.26820/reciamuc/7.(1).enero.2023.151-162).
- Na K, Oh B-C, Jung Y. Multifaceted role of CD14 in innate immunity and tissue homeostasis. *Cytokine Growth Factor Rev.* 2023;74:100–107. <http://dx.doi.org/10.1016/j.cytogfr.2023.08.008>.
- Nahon P, Cobat A. Human genetics of HCV infection phenotypes in the era of direct-acting antivirals. *Hum Genet.* 2020;139:855–863. <http://dx.doi.org/10.1007/s00439-020-02136-4>.
- Nichols-Vinueza DX, Delmonte OM, Bundy V, Bosticardo M, Zimmermann MT, Dsouza NR, et al. POLD1 Deficiency Reveals a Role for POLD1 in DNA Repair and T and B Cell Development. *J Clin Immunol.* 2021;41:270–273. <http://dx.doi.org/10.1007/s10875-020-00903-6>.
- Nicolas E, Golemis EA, Arora S. POLD1: Central mediator of DNA replication and repair, and implication in cancer and other pathologies. *Gene.* 2016;590:128–141. <http://dx.doi.org/10.1016/j.gene.2016.06.031>.
- Nielsen SCA, Boyd SD. Human adaptive immune receptor repertoire analysis—Past, present, and future. *Immunol Rev.* 2018;284:9–23. <http://dx.doi.org/10.1111/imr.12667>.
- O'Connor KS, Ahlenstiel G, Suppiah V, Schibeci S, Ong A, Leung R, et al. IFNL3 mediates interaction between innate immune cells: Implications for hepatitis C virus pathogenesis. *Innate Immun.* 2014;20:598–605. <http://dx.doi.org/10.1177/1753425913503385>.
- O'Neill LAJ, Golenbock D, Bowie AG. The history of Toll-like receptors — redefining innate immunity. *Nat Rev Immunol.* 2013;13:453–460. <http://dx.doi.org/10.1038/nri3446>.
- Oughtred R, Rust J, Chang C, Breitkreutz B, Stark C, Willems A, et al. The BIOGRID database: A comprehensive biomedical resource of curated protein, genetic, and chemical interactions. *Protein Sci.* 2021;30:187–200. <http://dx.doi.org/10.1002/pro.3978>.
- Paz y Miño Cepeda C. Rompecabezas evolutivo. Baltimore, Maryland: Project Muse; 2024.
- Paz-y-Miño Cepeda C, Morillo SA, Celi AP, Witte T, Muñoz Guijarro MJ, Collantes JC, et al. CCR5Δ, CCR2-64I, and SDF1-3'A Polymorphisms Related to Resistance to HIV-1 Infection and Disease in the Ecuadorian Population. *Hum Biol.* 2005;77:521–526. <http://dx.doi.org/10.1353/hub.2005.0068>.
- Paz-y-Miño C, Zambrano AK, Leone PE. Interactoma de predisposición y resistencia a SARS-CoV-2. *Proteínas, genes y funciones. Bionatura.* 2021;6:1555–1562. <http://dx.doi.org/10.21931/RB/2021.06.01.17>.
- Pecoraro V, Cuccorese M, Trenti T. Genetic polymorphisms of ACE1, ACE2, IFTM3, TMPRSS2 and TNFα genes associated with susceptibility and severity of SARS-CoV-2 infection: a systematic review and meta-analysis. *Clin Exp Med.* 2023;23:3251–3264. <http://dx.doi.org/10.1007/s10238-023-01038-9>.
- Peng B, Kong G, Yang C, Ming Y. Erythropoietin and its derivatives: from tissue protection to immune regulation. *Cell Death Dis.* 2020;11:79. <http://dx.doi.org/10.1038/s41419-020-2276-8>.
- Pleiman CM, D'Ambrosio D, Cambier JC. The B-cell antigen receptor complex: structure and signal transduction. *Immunol Today.* 1994;15:393–399. [http://dx.doi.org/10.1016/0167-5699\(94\)90267-4](http://dx.doi.org/10.1016/0167-5699(94)90267-4).
- Qi S, Sun C, Wang J, Wo L, Li Y, Wang C, et al. Identification of NECTIN1 as a novel restriction factor for flavivirus infection. Meng X-J, Weger-Lucarelli J, editors. *mBio.* 2024;15:e02708-24. <http://dx.doi.org/10.1128/mbio.02708-24>.
- Quintana-Murci L. Human Immunology through the Lens of Evolutionary Genetics. *Cell.* 2019;177:184–199. <http://dx.doi.org/10.1016/j.cell.2019.02.033>.
- Rajbhandari L, Shukla P, Jagdish B, Mandalla A, Li Q, Ali MA, et al. Nectin-1 Is an Entry Mediator for Varicella-Zoster Virus Infection of Human Neurons. Goodrum F, editor. *J Virol.* 2021;95:e01227-21. <http://dx.doi.org/10.1128/JVI.01227-21>.
- Ramezani A, Nahad MP, Faghiloo E. The role of Nrf2 transcription factor in viral infection. *J Cell Biochem.* 2018;119:6366–6382. <http://dx.doi.org/10.1002/jcb.26897>.
- Ramos-Casals M, Brahmer JR, Callahan MK, Flores-Chávez A, Keegan N, Khamashta MA, et al. Immune-related adverse events of checkpoint inhibitors. *Nat Rev Dis Primer.* 2020;6:38. <http://dx.doi.org/10.1038/s41572-020-0160-6>.
- Rankin JA. Biological Mediators of Acute Inflammation: AACN Clin Issues Adv Pract Acute Crit Care. 2004;15:3–17. <http://dx.doi.org/10.1097/00044067-200401000-00002>.
- Rasch RFR. Ancient History and New Frontiers: Infectious Diseases. *Nurs Clin North Am.* 2019;54:xv–xvi. <http://dx.doi.org/10.1016/j.cnur.2019.03.001>.
- Rast J, Litman G. Evolution of innate and adaptive immune recognition structures. *Semin Immunol.* 2010;22:1–2. <http://dx.doi.org/10.1016/j.smim.2010.01.001>.
- Redondo-García S, Barritt C, Papagregoriou C, Yeboah M, Frendeus B, Cragg MS, et al. Human leukocyte immunoglobulin-like receptors in health and disease. *Front Immunol.* 2023;14:1282874. <http://dx.doi.org/10.3389/fimmu.2023.1282874>.
- Rickert RC. New insights into pre-BCR and BCR signalling with relevance to B cell malignancies. *Nat Rev Immunol.* 2013;13:578–591. <http://dx.doi.org/10.1038/nri3487>.
- Ricklin D, Reis ES, Mastellos DC, Gros P, Lambris JD. Complement component C3 – The “Swiss Army Knife” of innate immunity and host defense. *Immunol Rev.* 2016;274:33–58. <http://dx.doi.org/10.1111/imr.12500>.
- Robledo D, Matika O, Hamilton A, Houston RD. Genome-Wide Association and Genomic Selection for Resistance to Amoebic Gill Disease in Atlantic Salmon. *G3 GenesGenomesGenetics.* 2018;8:1195–1203. <http://dx.doi.org/10.1534/g3.118.200075>.
- Rolland T, Taşan M, Charloreaux B, Pevzner SJ, Zhong Q, Sahni N, et al. A Proteome-Scale Map of the Human Interactome Network. *Cell.* 2014;159:1212–1226. <http://dx.doi.org/10.1016/j.cell.2014.10.050>.
- Rothfels K, Milacic M, Matthews L, Haw R, Sevilla C, Gillespie M, et al. Using the Reactome Database. *Curr Protoc.* 2023;3:e722. <http://dx.doi.org/10.1002/cpz1.722>.
- Rotival M. Characterising the genetic basis of immune response variation to identify causal mechanisms underlying disease susceptibility. *HLA.* 2019;94:275–284. <http://dx.doi.org/10.1111/tan.13598>.
- Saengsiwaritt W, Jittikoon J, Chaikledkaew U, Udomsinprasert W. Genetic polymorphisms of ACE1, ACE2, and TMPRSS2 associated with COVID-19 severity: A systematic review with meta-analysis. *Rev Med Virol.* 2022;32:e2323. <http://dx.doi.org/10.1002/rmv.2323>.
- Santiago VE, Dombrowski JG, Kawahara R, Rosa-Fernandes L, Mule SN, Murillo O, et al. Complement System Activation Is a Plasma Biomarker Signature during Malaria in Pregnancy. *Genes.* 2023;14:1624. <http://dx.doi.org/10.3390/genes14081624>.
- Setliff I, Shiakolas AR, Pilewski KA, Murji AA, Mapengo RE, Janowska K, et al. High-Throughput Mapping of B Cell Receptor Sequences to Antigen Specificity. *Cell.* 2019;179:1636–1646.e15. <http://dx.doi.org/10.1016/j.cell.2019.11.003>.
- Sharygin D, Koniaris LG, Wells C, Zimmers TA, Hamidi T. Role of CD14 in human disease. *Immunology.* 2023;169:260–270. <http://dx.doi.org/10.1111/imm.13634>.
- Shi Y, Su J, Chen R, Wei W, Yuan Z, Chen X, et al. The Role of Innate Immunity in Natural Elite Controllers of HIV-1 Infection. *Front Immunol.* 2022;13:780922. <http://dx.doi.org/10.3389/fimmu.2022.780922>.
- Silva AMVD, Alvarado-Arnez LE, Azamor T, Batista-Silva LR, Leal-Calvo T, Bezerra OCDL, et al. Interferon-lambda 3 and 4 Polymorphisms Increase Sustained Virological Responses and Regulate Innate Immunity in Antiviral Therapy With Pegylated Interferon-Alpha. *Front Cell Infect Microbiol.* 2021;11:656393. <http://dx.doi.org/10.3389/fcimb.2021.656393>.

- Smatti MK, Alkhatib HA, Al Thani AA, Yassine HM. Will Host Genetics Affect the Response to SARS-CoV-2 Vaccines? Historical Precedents. *Front Med*. 2022;9:802312. <http://dx.doi.org/10.3389/fmed.2022.802312>.
- Stefani C, Sangalli A, Locatelli E, Federico T, Malerba G, Romanelli MG, et al. Increased Prevalence of Unstable HLA-C Variants in HIV-1 Rapid-Progressor Patients. *Int J Mol Sci*. 2022;23:14852. <http://dx.doi.org/10.3390/ijms232314852>.
- Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, et al. The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence Analyses. *Curr Protoc Bioinforma* [Internet]. 2016 [cited 2024 Nov 5];54. <http://dx.doi.org/10.1002/cpbi.5>.
- Stertz S, Hale BG. Interferon system deficiencies exacerbating severe pandemic virus infections. *Trends Microbiol*. 2021;29:973–982. <http://dx.doi.org/10.1016/j.tim.2021.03.001>.
- Su N, Hu C, Shao T, Jin C, Li H, Ji J, et al. Functional role of CD40 and CD154 costimulatory signals in IgZ-mediated immunity against bacterial infection. *Fish Shellfish Immunol Rep*. 2021;2:100038. <http://dx.doi.org/10.1016/j.fsirep.2021.100038>.
- Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res*. 2023;51:D638–D646. <http://dx.doi.org/10.1093/nar/gkac1000>.
- Tanaka S, Baba Y. B Cell Receptor Signaling. In: Wang J-Y, editor. *B Cells Immun Toler* [Internet]. Singapore: Springer Singapore; 2020 [cited 2024 Nov 5]. p. 23–36. Available from: http://link.springer.com/10.1007/978-981-15-3532-1_2.
- Tarris G, De Rougemont A, Charkaoui M, Michiels C, Martin L, Belliot G. Enteric Viruses and Inflammatory Bowel Disease. *Viruses*. 2021;13:104. <http://dx.doi.org/10.3390/v13010104>.
- The UniProt Consortium, Bateman A, Martin M-J, Orchard S, Magrane M, Ahmad S, et al. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Res*. 2023;51:D523–D531. <http://dx.doi.org/10.1093/nar/gkac1052>.
- Vasconcelos RHT, Montenegro SML, Azevedo EAN, Gomes YM, Morais CNL. Genetic susceptibility to chronic Chagas disease: An overview of single nucleotide polymorphisms of cytokine genes. *Cytokine*. 2012;59:203–208. <http://dx.doi.org/10.1016/j.cyto.2012.04.035>.
- Vasseur E, Quintana-Murci L. The impact of natural selection on health and disease: uses of the population genetics approach in humans. *Evol Appl*. 2013;6:596–607. <http://dx.doi.org/10.1111/eva.12045>.
- Velastegui E, Vera E, Vanden Berghe W, Muñoz MS, Orellana-Manzano A. “HLA-C: evolution, epigenetics, and pathological implications in the major histocompatibility complex.” *Front Genet*. 2023;14:1206034. <http://dx.doi.org/10.3389/fgene.2023.1206034>.
- Velavan TP, Pallerla SR, Rüter J, Augustin Y, Kreamsner PG, Krishna S, et al. Host genetic factors determining COVID-19 susceptibility and severity. *eBioMedicine*. 2021;72:103629. <http://dx.doi.org/10.1016/j.ebiom.2021.103629>.
- Vijay K. Toll-like receptors in immunity and inflammatory diseases: Past, present, and future. *Int Immunopharmacol*. 2018;59:391–412. <http://dx.doi.org/10.1016/j.intimp.2018.03.002>.
- Villarino AV, Kanno Y, O’Shea JJ. Mechanisms and consequences of Jak–STAT signaling in the immune system. *Nat Immunol*. 2017;18:374–384. <http://dx.doi.org/10.1038/ni.3691>.
- Walker TM, Kohl TA, Omar SV, Hedge J, Del Ojo Elias C, Bradley P, et al. Whole-genome sequencing for prediction of Mycobacterium tuberculosis drug susceptibility and resistance: a retrospective cohort study. *Lancet Infect Dis*. 2015;15:1193–1202. [http://dx.doi.org/10.1016/S1473-3099\(15\)00062-6](http://dx.doi.org/10.1016/S1473-3099(15)00062-6).
- Wang H, Liu M. Complement C4, Infections, and Autoimmune Diseases. *Front Immunol*. 2021;12:694928. <http://dx.doi.org/10.3389/fimmu.2021.694928>.
- Wang J, Chai Q, Lei Z, Wang Y, He J, Ge P, et al. LILRB1-HLA-G axis defines a checkpoint driving natural killer cell exhaustion in tuberculosis. *EMBO Mol Med*. 2024;16:1755–1790. <http://dx.doi.org/10.1038/s44321-024-00106-1>.
- Wang J, Ganaie SS, Cheng F, Xu P, Ning K, Wang X, et al. RNA Binding Motif Protein RBM45 Regulates Expression of the 11-Kilodalton Protein of Parvovirus B19 through Binding to Novel Intron Splicing Enhancers. Meng X-J, editor. *mBio*. 2020;11:e00192-20. <http://dx.doi.org/10.1128/mBio.00192-20>.
- Wang Y, Qian G, Zhu L, Zhao Z, Liu Y, Han W, et al. HIV-1 Vif suppresses antiviral immunity by targeting STING. *Cell Mol Immunol*. 2022;19:108–121. <http://dx.doi.org/10.1038/s41423-021-00802-9>.
- Wang Y, Sun Q, Zhang Y, Li X, Liang Q, Guo R, et al. Systemic immune dysregulation in severe tuberculosis patients revealed by a single-cell transcriptome atlas. *J Infect*. 2023;86:421–438. <http://dx.doi.org/10.1016/j.jinf.2023.03.020>.
- Wang Y-T, Ji S-X, Xia X-Y, Wan T-J, Zou J-X, Li B-X, et al. Proteomic investigation and biomarker identification of lung and spleen deficiency syndrome in HIV/AIDS immunological nonresponders. *J Thorac Dis*. 2023;15:1460–1472. <http://dx.doi.org/10.21037/jtd-23-322>.
- Wen C, Dong Z, Wang Y, Ye G, Ma Y, Yi X, et al. CTLA4+CD4+CXCR5–FOXP3+ T cells associate with unfavorable outcome in patients with chronic HBV infection. *BMC Immunol*. 2023;24:3. <http://dx.doi.org/10.1186/s12865-022-00537-w>.
- Werner M, Schefczyk S, Trippler M, Treckmann JW, Baba HA, Gerken G, et al. Antiviral Toll-like Receptor Signaling in Non-Parenchymal Liver Cells Is Restricted to TLR3. *Viruses*. 2022;14:218. <http://dx.doi.org/10.3390/v14020218>.
- Wu J, Tan Y, Deng H, Zhu D. Relationship between degree–rank function and degree distribution of protein–protein interaction networks. *Comput Biol Chem*. 2008;32:1–4. <http://dx.doi.org/10.1016/j.compbiolchem.2007.08.008>.
- Wu Y-H, Chang W-T, Hsu C-L, Lu Y-F, Wang J-T, Tzeng S-J. FcγRIIB modulates splenic germinal center response against immune subversion during acute influenza A virus infection. *J Microbiol Immunol Infect*. 2024;57:64–75. <http://dx.doi.org/10.1016/j.jmii.2023.11.007>.
- Wu Z, Mehrabi Nasab E, Arora P, Athari SS. Study effect of probiotics and prebiotics on treatment of OVA-LPS-induced of allergic asthma inflammation and pneumonia by regulating the TLR4/NF-κB signaling pathway. *J Transl Med*. 2022;20:130. <http://dx.doi.org/10.1186/s12967-022-03337-3>.
- Wudhikarn K, Perales M-A. Infectious complications, immune reconstitution, and infection prophylaxis after CD19 chimeric antigen receptor T-cell therapy. *Bone Marrow Transplant*. 2022;57:1477–1488. <http://dx.doi.org/10.1038/s41409-022-01756-w>.
- Xu Z, Li D, Qu W, Yin Y, Qiao S, Zhu Y, et al. Card9 protects sepsis by regulating Ripk2-mediated activation of NLRP3 inflammasome in macrophages. *Cell Death Dis*. 2022;13:502. <http://dx.doi.org/10.1038/s41419-022-04938-y>.
- Yao P-A, Sun H-J, Li X-Y. Identification of key genes in late-onset major depressive disorder through a co-expression network module. *Front Genet*. 2022;13:1048761. <http://dx.doi.org/10.3389/fgene.2022.1048761>.
- Zhu M, Ma X, Huang J, Lu F, Chen Y, Hu J, et al. Extracellular vesicle-derived miR-1249–5p regulates influenza A virus-induced acute lung injury in RAW246.7 cells through targeting SLC4A1. *Microbes Infect*. 2022;24:104998. <http://dx.doi.org/10.1016/j.micinf.2022.104998>.